

① LEVEL II

AD A088661

DEPARTMENT OF STATISTICS

The Ohio State University

OSU

DTIC  
ELECTE  
SEP 3 1980  
S B D

COLUMBUS, OHIO

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

80 8 4 037

FILE COPY

① LEVEL II

ESTIMATION OF THE LARGER MEAN

by

Ishwari D. Dhariyal  
Edward J. Dudewicz\*  
Saul Blumenthal

DTIC  
ELECTE  
SEP 3 1980  
B

Technical Report No. 188  
Department of Statistics  
The Ohio State University  
Columbus, Ohio 43210  
July 1980

\* This author's research was supported by Office of Naval Research Contract  
No. N00014-78-C-0543.

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 188	2. GOVT ACCESSION NO. AD-A088 667	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ESTIMATION OF THE LARGER MEAN.		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Ishwari D. Dhariyal Edward J. Dudewicz Saul Blumenthal		8. CONTRACT OR GRANT NUMBER(s) N00014-78-C-0543
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics The Ohio State University Columbus, Ohio 43210		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 042-403
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217		12. REPORT DATE July 1980
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) TH-100		13. NUMBER OF PAGES i + 20
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) <div style="border: 1px solid black; padding: 5px; text-align: center;"> <b>DISTRIBUTION STATEMENT A</b>            Approved for public release;            Distribution Unlimited         </div>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Estimation, Larger Mean, Maximum Probability Estimator, Iterated Bias Elimination Estimator, Bias, Mean Squared Error, Ordered Parameters.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Estimation of the larger of two normal means $\mu_1$ and $\mu_2$ is studied. Two new estimators are studied for this problem, the maximum probability estimator and the iterated bias elimination estimators. If $ \mu_1 - \mu_2 $ is not close to zero (as evidenced by a strongly consistent estimator), a maximum probability estimator is seen to be best as regards both bias and mean-squared error.		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

70-100-1

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DOC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
PER LETTER		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL	and/or SPECIAL
A		

## Estimation of the Larger Mean

by

Ishwari D. Dhariyal, Indian Institute of Technology, Kanpur, U. P.

Edward J. Dudewicz, The Ohio State University, Columbus, Ohio

Saul Blumenthal, University of Illinois, Urbana, Illinois

### Abstract

In the present paper estimation of the larger of two normal means is studied. Two new estimators are added to the class of possible estimators of the larger normal mean, namely, the maximum probability estimator and the iterated bias elimination estimators. If the magnitude of the difference between the two population means is not close to zero (as evidenced by its strongly consistent estimator, the magnitude of the difference between the two sample means) a suitably chosen maximum probability estimator is seen to be best as regards both bias and mean-squared error.

### 1. Introduction

Let  $X_{11}, \dots, X_{1n}$  be a random sample of size  $n$  from a normal population with mean  $\mu_1$  and variance  $\sigma^2$ ,  $i = 1, \dots, k$  ( $\geq 2$ ). Let  $\mu_{[1]} \leq \dots \leq \mu_{[k]}$  be the ordered unknown means and suppose  $\sigma^2$  is known. The estimation analog of the well-known ranking and selection problem [see Bechhofer, Kiefer, and Sobel (1968)] has been the topic of inquiry by Alam (1967), Blumenthal (1975, 1976), Blumenthal and Cohen (1968), Dudewicz (1971, 1972, 1973, 1976) and others. Once the decision has been made as to which of the  $k$  populations has the largest mean,

it is natural to ask, "How large is this largest mean?" For two examples of application of estimation of the larger normal mean, the reader is referred to Blumenthal and Cohen (1968).

In this paper we study this problem and attempt to enlarge the class of estimators of the larger mean. In section 2 we investigate, mostly numerically, the behavior of a class of estimators called Maximum Probability Estimators (MPE's) introduced by Weiss and Wolfowitz (1967). Dudewicz (1973) first gave the MPE's of the ranked means. In section 3 a new class of estimators called Iterated Bias Elimination Estimators (IBEE's) are introduced and investigated. A comparison along the lines of Blumenthal and Cohen (1968) is made in section 4.

## 2. Maximum Probability Estimators

Definition (Weiss and Wolfowitz). Let  $\Theta$  be a closed region in the  $m$ -dimensional Euclidean space  $R^m$ ,  $\Theta \subset \bar{\Theta}$ , where  $\bar{\Theta}$  is a closed region such that every finite boundary point of  $\Theta$  is an inner point of  $\bar{\Theta}$ . For each  $n$  let  $X(n)$  denote the (finite) vector of random variables of which the estimator is to be a function. Let  $K_n(x, \theta)$  be the density of  $X(n)$  with respect to a sigma-finite measure. Let  $R$  be a fixed bounded set in  $R^m$  and let  $k(n) = (k_1(n), \dots, k_m(n))$  be a sequence of numbers such that  $k_i(n) \rightarrow \infty$  ( $n \rightarrow \infty$ ) for each  $i$ . Let  $d = (d_1, \dots, d_m)$  and  $d - R/k(n) = \{(z_1, \dots, z_m) \in \bar{\Theta} : d_i - y_i/k_i(n) = z_i, i = 1, \dots, m; (y_1, \dots, y_m) \in R\}$ . Then  $Z_n$  is an MPE with respect to  $R$  and  $k(n)$  if  $Z_n(x)$  equals a  $d \in \bar{\Theta}$  such that

$$\int_{d-R/k(n)} K_n(x, \theta) d\theta = \sup_{t \in \bar{\Theta}} \int_{t-R/k(n)} K_n(x, \theta) d\theta. \quad (2.1)$$

Now let  $\bar{X}_1, \dots, \bar{X}_k$  denote the sample means. Let in the above definition

$$X(n) = (\bar{X}_{[1]}, \dots, \bar{X}_{[k]}),$$

$$\theta \in \mathbb{R}^k,$$

$$\Theta = \{\underline{\mu} = (\mu_1, \dots, \mu_k) \in \mathbb{R}^k : \mu_1 = \mu_{[1]}, \dots, \mu_k = \mu_{[k]}\}$$

and

$$K_n(x, \theta) = f_{\bar{X}_{[1]}, \dots, \bar{X}_{[k]}}(x_1, \dots, x_k; \mu)$$

with respect to Lebesgue measure on  $\mathbb{R}^k$  where  $\bar{X}_{[1]} \leq \dots \leq \bar{X}_{[k]}$  are ordered sample means. (Note that  $K_n(x, \theta) > 0$  iff  $x_1 \leq \dots \leq x_k$ .) Let

$$k(n) = (\sqrt{n}/\sigma, \dots, \sqrt{n}/\sigma)$$

and

$$R = (-r_1/2, r_1/2) \times (-r_2/2, r_2/2) \times \dots \times (-r_k/2, r_k/2),$$

$r_1, \dots, r_k$  being positive real numbers. Define

$$d-R/k(n) = \left\{ (\mu_1, \dots, \mu_k) \in \mathbb{R}^k : d_i - \frac{\sigma r_1}{2\sqrt{n}} \leq \mu_i \leq d_i + \frac{\sigma r_1}{2\sqrt{n}}, i = 1, \dots, k \right\}.$$

We know that

$$f_{\bar{X}_{[1]}, \dots, \bar{X}_{[k]}}(x_1, \dots, x_k; \mu) = (\sqrt{n}/\sigma)^k \sum_{\beta \in S_k} \prod_{i=1}^k \phi\left(\frac{x_{\beta(i)} - \mu_i}{\sigma/\sqrt{n}}\right),$$

where  $S_k$  is the set of permutations on integers  $1, \dots, k$  and  $\phi(\cdot)$  denotes the standard normal density function. With  $d_{\beta(i)} = \sqrt{n}(x_{\beta(i)} - x_1)/\sigma$  and

$t_i = x_1 + a_i \sigma/\sqrt{n}$  we find from the definition that  $t = (t_1, \dots, t_k)$  is an MPE for  $\mu = (\mu_{[1]}, \dots, \mu_{[k]})$  if  $a = (a_1, \dots, a_k)$  are chosen so as to achieve

$$\sup_a \sum_{\beta \in S_k} \prod_{i=1}^k \left\{ \Phi(d_{\beta(i)} - a_i + r_1/2) - \Phi(d_{\beta(i)} - a_i - r_1/2) \right\} \quad (2.2)$$

where  $\Phi(\cdot)$  denotes the distribution function of a standard normal variable.

For values of  $k$  up to about 20 and using the observed values of  $d_1$ 's

we can find  $a_1, \dots, a_k$  by utilizing some function maximization (minimization) algorithm such as that of Nelder and Mead (1965).

In the rest of this section we investigate numerically the MPE's of  $(\mu_{[1]}, \mu_{[2]})$ , that is, for the case  $k = 2$ . For the values of the difference between the larger and the smaller sample means considered, after a preliminary analysis, we found that we could treat  $r_1, r_2 = 0.5, 1.5, \text{ and } 5.0$  as small, medium, and large respectively for the study. Thus we have nine pairs of  $(r_1, r_2)$  to look at. We found numerically that for  $d \leq \min(r_1, r_2)$ ,  $a_1 = -a_2 = d/2$  maximize

$$\begin{aligned} g(a_1, a_2) &= \sum_{\beta \in S_2} \prod_{i=1}^2 \left\{ \Phi(d_{\beta(i)} - a_1 + r_1/2) - \Phi(d_{\beta(i)} - a_1 - r_1/2) \right\} \\ &= \left\{ \Phi(a_1 + r_1/2) - \Phi(a_1 - r_1/2) \right\} \left\{ \Phi(a_2 + r_2/2) - \Phi(a_2 - r_2/2) \right\} \\ &\quad + \left\{ \Phi(a_1 - d + r_1/2) - \Phi(a_1 - d - r_1/2) \right\} \left\{ \Phi(a_2 + d + r_2/2) - \Phi(a_2 + d - r_2/2) \right\}, \end{aligned}$$

where  $d = \sqrt{n}(x_2 - x_1)/\sigma$ . This observation was checked with  $r_1, r_2$  changing in steps of 0.1 from 0.1 to 3.5. Redefining  $t_1$  and  $t_2$  as  $t_1 = x_1 + a_1\sigma/\sqrt{n}$  and  $t_2 = x_2 - a_2\sigma/\sqrt{n}$  we can write  $g(a_1, a_2)$  as

$$\begin{aligned} g(a_1, a_2) &= \left\{ \Phi(a_1 + r_1/2) - \Phi(a_1 - r_1/2) \right\} \left\{ \Phi(a_2 + r_2/2) - \Phi(a_2 - r_2/2) \right\} \\ &\quad + \left\{ \Phi(a_1 - d + r_1/2) - \Phi(a_1 - d - r_1/2) \right\} \left\{ \Phi(a_2 + d + r_2/2) - \Phi(a_2 + d - r_2/2) \right\} \quad (2.3) \end{aligned}$$

and now  $a_1 = a_2 = d/2$  maximize  $g(a_1, a_2)$  for  $d \leq \min(r_1, r_2)$ . Table 1 gives the values of  $(a_1, a_2)$  for the above nine pairs of  $(r_1, r_2)$  for  $d = 0.1(0.1)3.5$ . We tabulate for only six  $(r_1, r_2)$  pairs because of the subscript symmetry in (2.3) and hence the other three columns can be deduced from this table itself. For these calculations the Nelder-Mead (1965) simplex method was utilized. For

each of the nine  $(r_1, r_2)$  pairs and each value of  $d$ , it was verified that  $g(a_1^0 \pm 10^{-7}, a_2^0 \pm 10^{-7}) < g(a_1^0, a_2^0)$  where  $(a_1^0, a_2^0)$  is the calculated value. Thus the values reported are correct to seven decimal places.

Bias. Now without loss of generality, we consider the MPE's of  $\mu_{[2]}$  only. For given  $r_1, r_2$ , the MPE of  $\mu_{[2]}$  is given by

$$t_2 = t_2(r_1, r_2) = \bar{X}_{[2]} - \tau a_2(Z, r_1, r_2)$$

where  $\tau = \sigma/\sqrt{n}$  and  $Z = (\bar{X}_{[2]} - \bar{X}_{[1]})/2$ . Therefore, the bias is

$$\begin{aligned} B(t_2) &= B(t_2, w, \tau) \\ &= B(\bar{X}_{[2]}) - \tau \int_0^\infty a_2(z, r_1, r_2) f(z, w, \tau) dz \end{aligned} \quad (2.4)$$

where  $B(\bar{X}_{[2]})$  denotes the bias of  $\bar{X}_{[2]}$  as an estimator of  $\mu_{[2]}$  and

$$f(z, w, \tau) = \frac{\sqrt{2}}{\tau} \left\{ \phi(\sqrt{2}(z-w)/\tau) + \phi(\sqrt{2}(z+w)/\tau) \right\}, \quad z > 0 \quad (2.5)$$

is the density of  $Z$  with  $w = (\mu_{[2]} - \mu_{[1]})/2$ . Also from Blumenthal and Cohen (1968) we have

$$B(\bar{X}_{[2]}) = \frac{\tau}{\sqrt{\pi}} e^{-w^2/\tau^2} - 2w\phi(-\sqrt{2} w/\tau). \quad (2.6)$$

From (2.4), (2.5), and (2.6) it is clear that  $B(t_2, w, \tau) = \tau B(t_2, w/\tau, 1)$ . Therefore, we take  $\tau = 1$  in the following calculations and the values reported are in units of  $\tau$ . Values of  $B(\bar{X}_{[2]})$  are given in Blumenthal and Cohen (1968); we independently verified these values.

Now, from Dudewicz (1973) we know that  $0 < a_2 < 2Z$ . Also  $\phi(\sqrt{2}(z+w)) < \phi(\sqrt{2}(z-w))$ . Therefore, if we approximate  $\int_0^\infty a_2(z, r_1, r_2) f(z, w) dz$  by  $\int_0^M a_2(z, r_1, r_2) f(z, w) dz$ , the error due to this truncation is



$$E_T = \int_M^{\infty} a_2(z, r_1, r_2) f(z, \omega) dz$$

$$\leq 4\sqrt{2} \int_M^{\infty} z \phi(\sqrt{2}(z-\omega)) dz.$$

In order to bound  $E_T$  by  $\epsilon$ , it suffices to find an  $M = M_{\epsilon}(\omega)$  such that

$$4\sqrt{2} \int_{M_{\epsilon}(\omega)}^{\infty} z \phi(\sqrt{2}(z-\omega)) dz \leq \epsilon.$$

Prior to the numerical evaluation of the integral  $I_M = \int_0^{M_{\epsilon}(\omega)} a_2(z, r_1, r_2) f(z, \omega) dz$  a study of the function  $g(z, r_1, r_2, \omega) = a_2(z, r_1, r_2) f(z, \omega)$  revealed that this function has a sharp peak (possibly a non-differentiable point) in the interval  $[0, M_{\epsilon}(\omega)]$ . Three typical functions are graphed in Figure 1. This fact is an indication that one should evaluate the integral  $I_M$  in two parts, namely, in intervals  $[0, a]$  and  $[a, b]$ , where  $a = a(\omega)$  is the point such that  $a_2(z, r_1, r_2) = z$  for  $z \leq a$  [note that  $a$  turns out to be the same point where  $a_2(z, r_1, r_2)$  starts decreasing] and  $b = b(\omega)$  is such that  $a_2(z, r_1, r_2) \leq \epsilon$  for  $z \geq b$ . It was found that in each case considered  $b \leq M_{\epsilon}(\omega)$ . Since for  $z \geq b$ ,  $a_2(z, r_1, r_2) \leq \epsilon$ .

$$\int_b^{\infty} a_2(z, r_1, r_2) f(z, \omega) dz \leq \epsilon.$$

Hence we used Gaussian quadrature formula [see Stroud and Secrest (1966)]

$$\int_0^a a_2(z, r_1, r_2) f(z, \omega) dz + \int_a^b a_2(z, r_1, r_2) f(z, \omega) dz$$

to approximate

$$\int_0^{\infty} a_2(z, r_1, r_2) f(z, \omega) dz.$$

To control the error due to quadrature, for each  $(r_1, r_2)$  pair each of the integrals

$$I_1 = \int_0^a a_2(z, r_1, r_2) f(z, w) dz$$

and

$$I_2 = \int_a^b a_2(z, r_1, r_2) f(z, w) dz$$

was evaluated in 3, 5, 7, or 9 subintervals using 64 and 128 point Gaussian quadrature formulas. The criterion used to stop subdividing each subinterval was that the two values of the integral  $I_i$  obtained using 64 and 128 points respectively for a subdivision of the subinterval differ by no more than  $\epsilon$ ,  $i = 1, 2$ . Thus our approximation involves an error due to quadrature plus an error due to truncation which is bounded above by  $\epsilon + \epsilon^*$ . We used  $\epsilon = \epsilon^* = 10^{-7}$ . Thus we have

$$B(t_2) \approx B(\bar{X}_{[2]}) - I_1 - I_2$$

Mean-squared error (MSE). We have

$$\begin{aligned} \text{MSE}(t_2) &= \text{MSE}(t_2, w, \tau) \\ &= \text{MSE}(\bar{X}_{[2]}) + E [\tau^2 a_2^2(Z, r_1, r_2) - 2\tau(Z-w)a_2(Z, r_1, r_2)] \\ &= \text{MSE}(\bar{X}_{[2]}) - \tau^2 \int_0^w \left\{ 2\left(\frac{Z}{\tau} - \frac{w}{\tau}\right) a_2(Z, r_1, r_2) - a_2^2(Z, r_1, r_2) \right\} f(Z, w, \tau) dZ. \end{aligned} \quad (2.7)$$

Also

$$\text{MSE}(\bar{X}_{[2]}) = \tau^2 + w^2 \Phi(-\sqrt{2} w/\tau) - (2w\tau/\sqrt{\pi}) e^{-w^2/\tau^2} \quad (2.8)$$

has been tabulated by Blumenthal and Cohen (1968); we independently verified these values. Again, it is clear that  $\text{MSE}(t_2, w, \tau) = \text{MSE}(t_2, \frac{w}{\tau}, 1)$ . All the

calculations run parallel to those for the bias except that new bounds are found to approximate the integral in (2.7).

Tables 2 and 3 below give the bias and MSE respectively of  $t_2(r_1, r_2)$ . We see that smaller values of  $r_1, r_2$  give rise to smaller |bias| and smaller MSE.

### 3. Iterated Bias Elimination Estimators

Consider the case of  $k = 2$  normal populations with means  $\mu_1$  and  $\mu_2$  and a common known variance  $\sigma^2$ . Let  $X^{*(1)} = \bar{X}_{[2]}$  and let  $B^{*(1)}(\omega)$  denote the bias of  $X^{*(1)}$  as an estimator of  $\mu_{[2]}$ . Let  $\hat{B}^{*(1)}(\omega)$  denote a consistent estimator of  $B^{*(1)}(\omega)$ . Define  $X^{*(2)} = X^{*(1)} - \hat{B}^{*(1)}(\omega)$  which has a bias of  $B^{*(2)}(\omega)$  as an estimator of  $\mu_{[2]}$ . Let  $\hat{B}^{*(2)}(\omega)$  denote a consistent estimator of  $B^{*(2)}(\omega)$ . Continuing the process, let  $X^{*(m)} = X^{*(m-1)} - \hat{B}^{*(m-1)}(\omega)$ ,  $m = 2, 3, \dots$ . In this section we investigate the class of estimators  $\{X^{*(m)}; m = 1, 2, \dots\}$  of  $\mu_{[2]}$ .

Let  $C^{*(0)}(\omega) = \frac{\tau}{\sqrt{\pi}} e^{-\omega^2/\tau^2} - 2\omega\tau(\sqrt{2}\omega/\tau)$ , and  $C^{*(m)}(\omega) = \frac{\sqrt{m+1}}{\sqrt{\pi}} \tau e^{-\omega^2/\tau^2(m+1)} - 2\omega\tau(\sqrt{2}\omega/\sqrt{m+1}\tau)$ ,  $m = 1, 2, \dots$ . We estimate  $\omega$  (which appears in  $B^{*(m)}(\omega)$  through the continuous function  $C^{*(m-1)}(\omega)$ ,  $m = 1, 2, \dots$ ) by its strongly consistent estimator  $Z$ . That is,  $\hat{B}^{*(m)}(\omega) = B^{*(m)}(Z)$ .

Definition.  $X^{*(m)}$  is called the  $m$ -th IBEE of  $\mu_{[2]}$ ,  $m = 1, 2, \dots$ .

Lemma. For  $m = 1, 2, \dots$

$$E C^{*(m-1)}(Z) = C^{*(m)}(\omega) - C^{*(0)}(\omega).$$

Proof. The proof is straightforward involving just the routine evaluation of integrals of type

$$\int_0^{\infty} e^{-a_1 z^2} \phi(b_1 z + c_1) dz + \int_0^{\infty} z \phi(a_2 z) \phi(b_2 z + c_2) dz.$$

Theorem. For  $m = 1, 2, \dots$

$$B^{*(m)}(\omega) = \sum_{r=0}^{m-1} (-1)^r \binom{m}{r+1} C^{*(r)}(\omega).$$

Proof. From Blumenthal and Cohen (1968) we know that the theorem is true for  $m = 1$ . Suppose it is true for  $m = k$ . Then, by definition

$$X^{*(k+1)} = X^{*(k)} - \sum_{r=0}^{k-1} (-1)^r \binom{k}{r+1} C^{*(r)}(Z).$$

Therefore,

$$\begin{aligned} B^{*(k+1)}(\omega) &= B^{*(k)}(\omega) - \sum_{r=0}^{k-1} (-1)^r \binom{k}{r+1} E C^{*(r)}(Z) \\ &= \sum_{r=0}^{k-1} (-1)^r \binom{k}{r+1} C^{*(r)}(\omega) - \sum_{r=0}^{k-1} (-1)^r \binom{k}{r+1} \{C^{*(r+1)}(\omega) - C^{*(0)}(\omega)\} \\ &= \sum_{r=0}^k (-1)^r \binom{k+1}{r+1} C^{*(r)}(\omega). \end{aligned}$$

Thus, the theorem is true for  $m = k+1$ . The proof now follows by induction.

Corollary. For  $m = 2, 3, \dots$

$$X^{*(m)} = X^{*(1)} - \sum_{r=0}^{m-2} (-1)^r \binom{m}{r+2} C^{*(r)}(Z).$$

A numerical study as to the behavior of  $B^{*(m)}(\omega)$  as  $m$  increases shows that for smaller values of  $\omega$   $B^{*(m)}(\omega)$  decreases as  $m$  is increased. The behavior is reversed for larger values of  $\omega$ . Table 4 gives for  $m = 1, 2, \dots, 30$

$$b(m) = \max_{w=0.0(0.1)5.0} |B^{*(m)}(w)|$$

which reveals that for values of  $m$  and  $w$  considered  $X^{*(18)}$  is the minimax  $|bias|$  IBEE.

#### 4. Comparison of Estimators of $\mu_{[2]}$

In this section we make comparisons between the "best" estimators of Blumenthal and Cohen (1968), the "best" MPE and the "minimax"  $|bias|$  IBEE as regards their bias and MSE. From tables and graphs of Blumenthal and Cohen we find that with respect to the bias the estimator

$$\delta_H(1) = \begin{cases} (\bar{X}_1 + \bar{X}_2)/2 & \text{if } z < \tau \\ \bar{X}_{[2]} & \text{if } z \geq \tau \end{cases}$$

seems to be the "best" among all the estimators considered and with respect to MSE, the estimator

$$\delta_P = \frac{\bar{X}_1 + \bar{X}_2}{2} + zZ[\Phi(z/\sqrt{2}\tau) - \frac{1}{2}] + \frac{\tau}{\sqrt{\pi}} e^{-z^2/\tau^2}$$

seems to have an advantage over the other estimators. From section 2 we know that among the nine MPE's considered  $t_2(0.5, 0.5)$  is the 'best' both in case of bias and MSE. In Tables 5 and 6 we give the bias and MSE respectively of the four estimators  $\delta_H(1)$ ,  $\delta_P$ ,  $t_2(0.5, 0.5)$ , and  $X^{*(18)}$  and compare them graphically in Figures 2 and 3. (Note that the  $MSE(X^{*(18)})$  was calculated using Monte Carlo techniques.)

#### ACKNOWLEDGMENTS

This research was supported by Office of Naval Research Contract No. N00014-78-C-0543.



Table 2: Bias of Maximum Probability Estimators

n	r <sub>1</sub> =0.5		r <sub>1</sub> =0.5		r <sub>1</sub> =0.5		r <sub>1</sub> =1.5		r <sub>1</sub> =1.5		r <sub>1</sub> =5.0		r <sub>1</sub> =5.0		r <sub>1</sub> =5.0		r <sub>1</sub> =5.0	
	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5	r <sub>2</sub> =0.5	r <sub>2</sub> =1.5
0.0	.30462	.28916	.16334	.28950	.28950	.27224	.13703	.13703	.18708	.15778	.01186	.01186	.18708	.15778	.01186	.01186	.18708	.15778
0.1	.21134	.19583	.07088	.19618	.19618	.17885	.04212	.04212	.09391	.06330	-.08710	-.08710	.09391	.06330	-.08710	-.08710	.09391	.06330
0.2	.13139	.11575	-.01249	.11610	.11610	.09856	-.04257	-.04257	.01166	-.02014	-.18338	-.18338	.01166	-.02014	-.18338	-.18338	.01166	-.02014
0.3	.06443	.04859	-.08478	.04895	.04895	.03110	-.11696	-.11696	-.05882	-.09250	-.27816	-.27816	-.05882	-.09250	-.27816	-.27816	-.05882	-.09250
0.4	.00991	.00616	-.14599	.00579	.00579	-.02400	-.18092	-.18092	-.11768	-.15377	-.36945	-.36945	-.11768	-.15377	-.36945	-.36945	-.11768	-.15377
0.5	-.03204	.04921	-.19622	-.04882	-.04882	-.06738	-.23434	-.23434	-.16516	-.20400	-.45705	-.45705	-.16516	-.20400	-.45705	-.45705	-.16516	-.20400
0.6	-.06506	.08144	-.23563	.08104	.08104	-.09987	-.27718	-.27718	-.20160	-.24333	-.54005	-.54005	-.20160	-.24333	-.54005	-.54005	-.20160	-.24333
0.7	-.08753	.10389	-.26455	.10348	.10348	-.12244	-.30951	-.30951	-.22753	-.27202	-.61741	-.61741	-.22753	-.27202	-.61741	-.61741	-.22753	-.27202
0.8	-.10156	.11774	-.28346	.11732	.11732	-.13622	-.33158	-.33158	-.24364	-.29055	-.68794	-.68794	-.24364	-.29055	-.68794	-.68794	-.24364	-.29055
0.9	-.10844	.12423	-.29203	.12382	.12382	-.14240	-.34384	-.34384	-.25080	-.29958	-.75037	-.75037	-.25080	-.29958	-.75037	-.75037	-.25080	-.29958
1.0	-.10946	.12466	-.29413	.12425	.12425	-.14228	-.34694	-.34694	-.25005	-.29998	-.80342	-.80342	-.25005	-.29998	-.80342	-.80342	-.25005	-.29998
1.1	-.10592	.12032	-.28780	.11992	.11992	-.13714	-.34179	-.34179	-.24256	-.29281	-.84587	-.84587	-.24256	-.29281	-.84587	-.84587	-.24256	-.29281
1.2	-.09902	.11245	-.27523	.11297	.11297	-.12823	-.32946	-.32946	-.22961	-.27928	-.87663	-.87663	-.22961	-.27928	-.87663	-.87663	-.22961	-.27928
1.3	-.08988	.10218	-.25769	.10182	.10182	-.11674	-.31119	-.31119	-.21252	-.26073	-.89437	-.89437	-.21252	-.26073	-.89437	-.89437	-.21252	-.26073
1.4	-.07946	.09053	-.23649	.09020	.09020	-.10373	-.28833	-.28833	-.19258	-.23851	-.90007	-.90007	-.19258	-.23851	-.90007	-.90007	-.19258	-.23851
1.5	-.06856	.07835	-.21291	.07805	.07805	-.09010	-.26222	-.26222	-.17102	-.21395	-.89214	-.89214	-.17102	-.21395	-.89214	-.89214	-.17102	-.21395
1.6	-.05784	.06633	-.18816	.06607	.06607	-.07650	-.23420	-.23420	-.14893	-.18831	-.87141	-.87141	-.14893	-.18831	-.87141	-.87141	-.14893	-.18831
1.7	-.04775	.05498	-.16329	.05475	.05475	-.06378	-.20550	-.20550	-.12724	-.16267	-.83869	-.83869	-.12724	-.16267	-.83869	-.83869	-.12724	-.16267
1.8	-.03860	.04466	-.13921	.04445	.04445	-.05206	-.17719	-.17719	-.10660	-.13797	-.79524	-.79524	-.10660	-.13797	-.79524	-.79524	-.10660	-.13797
1.9	-.03059	.03555	-.11660	.03538	.03538	-.04167	-.15015	-.15015	-.08782	-.11491	-.74272	-.74272	-.08782	-.11491	-.74272	-.74272	-.08782	-.11491
2.0	-.02376	.02776	-.09598	.02762	.02762	-.03272	-.12508	-.12508	-.07097	-.09400	-.68310	-.68310	-.07097	-.09400	-.68310	-.68310	-.07097	-.09400
2.1	-.01810	.02196	-.07765	.02115	.02115	-.02521	-.10242	-.10242	-.05633	-.07552	-.61855	-.61855	-.05633	-.07552	-.61855	-.61855	-.05633	-.07552
2.2	-.01353	.01598	-.06175	.01589	.01589	-.01906	-.08246	-.08246	-.04390	-.05961	-.55130	-.55130	-.04390	-.05961	-.55130	-.55130	-.04390	-.05961
2.3	-.00992	.01179	-.04827	.01172	.01172	-.01415	-.06527	-.06527	-.03361	-.04621	-.48353	-.48353	-.03361	-.04621	-.48353	-.48353	-.03361	-.04621
2.4	-.00714	.00854	-.03710	.00848	.00848	-.01031	-.05080	-.05080	-.02528	-.03520	-.41724	-.41724	-.02528	-.03520	-.41724	-.41724	-.02528	-.03520
2.5	-.00504	.00607	-.02503	.00603	.00603	-.00738	-.03887	-.03887	-.01867	-.02634	-.35413	-.35413	-.01867	-.02634	-.35413	-.35413	-.01867	-.02634
2.6	-.00350	.00423	-.02082	.00420	.00420	-.00518	-.02925	-.02925	-.01355	-.01937	-.29559	-.29559	-.01355	-.01937	-.29559	-.29559	-.01355	-.01937
2.7	-.00238	.00290	-.01521	.00288	.00288	-.00358	-.02164	-.02164	-.00966	-.01399	-.24258	-.24258	-.00966	-.01399	-.24258	-.24258	-.00966	-.01399
2.8	-.00159	.00195	-.01092	.00194	.00194	-.00242	-.01574	-.01574	-.00677	-.00993	-.19570	-.19570	-.00677	-.00993	-.19570	-.19570	-.00677	-.00993
2.9	-.00105	.00129	-.00771	.00128	.00128	-.00161	-.01126	-.01126	-.00466	-.00693	-.15518	-.15518	-.00466	-.00693	-.15518	-.15518	-.00466	-.00693
3.0	-.00067	.00084	-.00535	.00083	.00083	-.00106	-.00792	-.00792	-.00315	-.00475	-.12092	-.12092	-.00315	-.00475	-.12092	-.12092	-.00315	-.00475
4.0	-.00000	.00000	-.00006	.00000	.00000	-.00001	-.00009	-.00009	-.00002	-.00004	-.00373	-.00373	-.00002	-.00004	-.00373	-.00373	-.00002	-.00004
5.0	-.00000	.00000	-.00000	.00000	.00000	-.00000	-.00000	-.00000	-.00000	-.00000	-.00002	-.00002	-.00000	-.00000	-.00002	-.00002	-.00000	-.00000

Table 3: MSE of Maximum Probability Estimators

	$r_1=0.5$ $r_2=0.5$	$r_1=0.5$ $r_2=1.5$	$r_1=0.5$ $r_2=5.0$	$r_1=1.5$ $r_2=0.5$	$r_1=1.5$ $r_2=1.5$	$r_1=1.5$ $r_2=5.0$	$r_1=5.0$ $r_2=0.5$	$r_1=5.0$ $r_2=1.5$	$r_1=5.0$ $r_2=5.0$
0.0	0.85293	0.83940	0.70630	0.83990	0.82437	0.67579	0.71356	0.71164	0.52502
0.1	0.81090	0.80040	0.69063	0.80084	0.78868	0.66526	0.72401	0.69769	0.53470
0.2	0.80138	0.79383	0.70551	0.79421	0.78531	0.68458	0.73594	0.71472	0.56785
0.3	0.81665	0.81204	0.74476	0.81236	0.80671	0.72816	0.77267	0.75654	0.62370
0.4	0.84949	0.84785	0.80252	0.84811	0.84577	0.79064	0.82782	0.81724	0.70154
0.5	0.89334	0.89472	0.87318	0.89491	0.89509	0.86684	0.89536	0.89113	0.80054
0.6	0.94245	0.94688	0.95145	0.94701	0.95158	0.91171	0.96965	0.97277	0.91957
0.7	0.99200	0.99944	1.03239	0.99950	1.00759	1.04041	1.04555	1.05702	1.05702
0.8	1.03818	1.04851	1.11150	1.04850	1.06003	1.12835	1.11850	1.13918	1.21063
0.9	1.07818	1.09118	1.18482	1.09108	1.10583	1.21133	1.18466	1.2510	1.37737
1.0	1.11023	1.12552	1.24910	1.12535	1.14295	1.28572	1.24102	1.28129	1.55337
1.1	1.13344	1.15057	1.30186	1.15033	1.17028	1.34854	1.28546	1.33515	1.73396
1.2	1.14776	1.16619	1.34152	1.16580	1.18757	1.39766	1.31680	1.37497	1.91370
1.3	1.15379	1.17292	1.39737	1.17257	1.19520	1.45185	1.33479	1.40002	2.08664
1.4	1.15260	1.17183	1.37955	1.17145	1.19450	1.45081	1.34004	1.41050	2.24655
1.5	1.14557	1.16435	1.37901	1.16395	1.18665	1.45509	1.33383	1.40745	2.38731
1.6	1.13421	1.15205	1.36729	1.15165	1.17339	1.44605	1.31799	1.39260	2.50331
1.7	1.12003	1.13654	1.34638	1.13614	1.15612	1.42562	1.29467	1.36816	2.58987
1.8	1.10439	1.11928	1.31854	1.11891	1.13734	1.39614	1.26613	1.33660	2.64362
1.9	1.08846	1.10157	1.28608	1.10122	1.11757	1.36016	1.23460	1.30045	2.66280
2.0	1.07312	1.08439	1.25118	1.08408	1.09825	1.32020	1.20205	1.26208	2.64740
2.1	1.05903	1.06851	1.21580	1.06823	1.08022	1.27861	1.17016	1.22359	2.59921
2.2	1.04659	1.05437	1.18155	1.05413	1.06406	1.23740	1.14021	1.18667	2.52163
2.3	1.03596	1.04221	1.14963	1.04201	1.05005	1.19821	1.11311	1.15250	2.41946
2.4	1.02716	1.03208	1.12087	1.03192	1.03828	1.16223	1.08937	1.12218	2.29841
2.5	1.02009	1.02387	1.09574	1.02374	1.02867	1.13022	1.06918	1.09586	2.16475
2.6	1.01455	1.01740	1.07438	1.01730	1.02104	1.10250	1.05248	1.07371	2.02473
2.7	1.01032	1.01243	1.05669	1.01235	1.01514	1.07917	1.03903	1.05557	1.88424
2.8	1.00718	1.00870	1.04240	1.00864	1.01067	1.06901	1.02846	1.04108	1.74837
2.9	1.00489	1.00597	1.03113	1.00593	1.00738	1.04465	1.02036	1.02978	1.62124
3.0	1.00327	1.00402	1.02244	1.00399	1.00500	1.03262	1.01428	1.02118	1.50582
4.0	1.00002	1.00003	1.00032	1.00003	1.00004	1.00053	1.00015	1.00025	1.02231
5.0	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00000	1.00014



Table 4: Values of  $b(m) = \max_{\theta=0.0(0.1)5.0} |B^{*(m)}(\theta)|$ , the

maximum |bias| of m-th IBEE.

m	b(m)	m	b(m)	m	b(m)
1	0.5642	2	0.3305	3	0.2761
4	0.2499	5	0.2338	6	0.2227
7	0.2144	8	0.2079	9	0.2027
10	0.1983	11	0.1945	12	0.1913
13	0.1885	14	0.1860	15	0.1832
16	0.1840	17	0.1822	18	0.1536
19	0.1879	20	0.2736	21	0.2047
22	1.4915	23	0.3366	24	4.3284
25	1.0571	26	16.4563	27	1.1254
28	105.9889	29	23.4225	30	109.0409

Table 5: Bias Comparison

$\mu$	$\delta_H(1)$	$\delta_P$	$t_2(0.5, 0.5)$	$\chi^*(18)$
0.0	0.2076	0.7979	0.3046	0.1536
0.1	0.1138	0.7019	0.2113	0.0669
0.2	0.0324	0.6138	0.1314	0.0016
0.3	-0.0367	0.5335	0.0644	-0.0389
0.4	-0.0938	0.4609	0.0099	-0.0655
0.5	-0.1392	0.3956	-0.0329	-0.0734
0.6	-0.1734	0.3374	-0.0651	-0.0746
0.7	-0.1972	0.2858	-0.0875	-0.0621
0.8	-0.2114	0.2404	-0.1016	-0.0446
0.9	-0.2169	0.2009	-0.1084	-0.0280
1.0	-0.2151	0.1667	-0.1095	-0.0219
1.1	-0.2071	0.1372	-0.1059	-0.0062
1.2	-0.1942	0.1122	-0.0990	0.0016
1.3	-0.1779	0.0911	-0.0899	0.0126
1.4	-0.1593	0.0733	-0.0795	0.0016
1.5	-0.1397	0.0586	-0.0686	0.0187
1.6	-0.1200	0.0465	-0.0578	0.0038
1.7	-0.1010	0.0366	-0.0478	0.0059
1.8	-0.0833	0.0286	-0.0386	0.0004
1.9	-0.0674	0.0221	-0.0306	-0.0010
2.0	-0.0535	0.0170	-0.0238	-0.0109
2.1	-0.0417	0.0129	-0.0181	-0.0057
2.2	-0.0318	0.0098	-0.0135	-0.0026
2.3	-0.0238	0.0073	-0.0100	-0.0228
2.4	-0.0175	0.0054	-0.0071	-0.0177
2.5	-0.0125	0.0040	-0.0050	-0.0104
2.6	-0.0089	0.0029	-0.0035	-0.0003
2.7	-0.0062	0.0021	-0.0024	-0.0076
2.8	-0.0042	0.0015	-0.0016	-0.0139
2.9	-0.0028	0.0011	-0.0011	-0.0024
3.0	-0.0019	0.0008	-0.0007	-0.0174
4.0	-0.0000	0.0000	-0.0000	-0.0015
5.0	0.0000	0.0000	-0.0000	0.0004

Table 6: MSE Comparison

$\omega$	$\delta_H(1)$	$\delta_P$	$t_2(0.5, 0.5)$	$x^*(18)$
0.0	0.7862	1.2182	0.8529	3.0538
0.1	0.7633	1.0455	0.8109	3.0725
0.2	0.7729	0.9067	0.8014	3.0582
0.3	0.8076	0.7970	0.8167	2.9850
0.4	0.8609	0.7113	0.8495	2.9157
0.5	0.9261	0.6453	0.8933	2.7558
0.6	0.9974	0.5989	0.9425	2.6158
0.7	1.0694	0.5674	0.9920	2.4251
0.8	1.1375	0.5520	1.0382	2.2187
0.9	1.1980	0.5496	1.0782	2.0066
1.0	1.2479	0.5590	1.1102	1.8548
1.1	1.2854	0.5795	1.1334	1.7210
1.2	1.3098	0.6070	1.1478	1.5996
1.3	1.3212	0.6417	1.1538	1.4964
1.4	1.3205	0.6794	1.1526	1.4021
1.5	1.3093	0.7189	1.1456	1.2870
1.6	1.2897	0.7583	1.1342	1.1914
1.7	1.2639	0.7956	1.1200	1.1325
1.8	1.2342	0.8305	1.1044	1.0869
1.9	1.2028	0.8615	1.0885	1.0673
2.0	1.1715	0.8888	1.0731	1.0432
2.1	1.1417	0.9120	1.0590	1.0229
2.2	1.1144	0.9315	1.0466	1.0163
2.3	1.0904	0.9474	1.0360	1.0130
2.4	1.0699	0.9601	1.0272	1.0109
2.5	1.0529	0.9702	1.0201	1.0194
2.6	1.0391	0.9780	1.0146	1.0207
2.7	1.0284	0.9840	1.0103	1.0213
2.8	1.0202	0.9885	1.0072	1.0206
2.9	1.0140	0.9918	1.0049	1.0185
3.0	1.0096	0.9942	1.0033	1.0155
4.0	1.0001	0.9999	1.0000	0.9979
5.0	1.0000	1.0000	1.0000	1.0024






Figure 1: Graphical investigation of behavior of MPE's.  $f$ , the density of  $Z$ ;  $a_2$ , the adjustment in  $\bar{X}_{[2]}$  given by MPE of  $u_{[2]}$ ; and  $g = a_2 f$  are plotted against  $Z$ . For all  $(r_1, r_2)$  pairs considered, (a), (b), (c) represent the typical plots when  $w = 3.0, 1.0, 0.0$  respectively.




Figure 2: Comparison of estimators of  $\mu_{[2]}$  with respect to their bias.




Figure 3: Comparison of estimators of  $\mu_{[2]}$  with respect to their MSE.

### References

- Alam, K. (1967): "A two-sample estimate of the largest mean," Annals of the Institute of Statistical Mathematics, 19, 271-283.
- Bechhofer, R. E., Kiefer, J., and Sobel, M. (1968): Sequential Identification and Ranking Procedures (with special reference to Koopman-Darmois populations), The University of Chicago Press, Chicago, Illinois.
- Blumenthal, S. (1975): "Sequential estimation of the largest normal mean when variance is unknown," Communications in Statistics, 4, 655-669.
- (1976): "Sequential estimation of the largest normal mean when variance is known," Annals of Statistics, 4, 1077-1087.
- , and Cohen, A. (1968): "Estimation of the larger of two normal means," Journal of the American Statistical Association, 63, 861-876.
- Dudewicz, E. J. (1971): "Maximum likelihood estimators for ranked means," Z. Wahrscheinlichkeitstheorie verw. Geb., 19, 29-42.
- (1972): "Point estimation of ordered parameters: the general location parameter case," Tamkang Journal of Mathematics, 3, 101-114.
- (1973): "Maximum probability estimators of ranked means," Annals of the Institute of Statistical Mathematics, 25, 467-477.
- (1976): "Estimation of ranked parameters: Insights and advances," Technical Report No. 118, Department of Statistics, Stanford University, Stanford, California.
- Nelder, J. A. and Mead, R. (1965): "A simplex method for function minimization," Computer Journal, 7, 308-313.
- Stroud, A. H. and Secrest, D. (1966): Gaussian Quadrature formulas, Prentice-Hall, Inc., Englewood Cliffs, New Jersey.
- Weiss, L. and Wolfowitz, J. (1967): "Maximum probability estimators," Annals of the Institute of Statistical Mathematics, 19, 193-206.